

211

**DROUGHT AND ACIDIFICATION OF A
MINEROTROPHIC CONIFER SWAMP ON
THE PRECAMBRIAN SHIELD, CANADA**

NOVEMBER 1992



Ontario

**Environment
Environnement**

22/10/92

ISBN 0-7778-0227-9

DROUGHT AND ACIDIFICATION OF A
MINEROTROPHIC CONIFER SWAMP
ON THE PRECAMBRIAN SHIELD, CANADA

NOVEMBER 1992



Cette publication technique
n'est disponible qu'en anglais.

Copyright: Queen's Printer for Ontario, 1992
This publication may be reproduced for non-commercial purposes
with appropriate attribution.

PIBS 2175

DROUGHT AND ACIDIFICATION OF A
MINEROTROPHIC CONIFER SWAMP
ON THE PRECAMBRIAN SHIELD, CANADA

Report prepared by:

Bruce D. LaZerte
Dorset Research Centre
Ontario Ministry of the Environment
P.O. Box 39
Dorset, Ontario P0A 1E0

NOVEMBER 1992

PIBS 2175

ABSTRACT

A seven year budget of a minerotrophic conifer swamp subjected to anthropogenically acidified deposition is presented. Contrary to other published studies, this swamp retains sulphate during wet years and exports it during dry years. Base cations are almost always exported. It is predicted that if this pattern continues, the cation exchange sites of the swamp will become increasingly saturated with protons and aluminum ions, and the waters passing through the swamp will be increasingly acidified.

TABLE OF CONTENTS

Abstract	i
Table of Contents	ii
List of Tables	iii
List of Figures	iv
Introduction	1
Methods	2
Errors of Method Checks	9
Results	11
Discussion	14
References	15
Appendix - Alkalinity Method	25

LIST OF TABLES

Table 1a	Quarterly and annual inputs, outputs and retention of water discharge (m^3), major ions ($\text{meq m}^{-2} \text{ quarter}^{-1}$) and DOC ($\text{g m}^{-2} \text{ quarter}^{-1}$) in Plastic 1 Swamp using Pc 1-08 data for un-channelized fluxes.
Table 1b	Quarterly and annual inputs, outputs and retention of major alkalinity components ($\text{meq m}^{-2} \text{ quarter}^{-1}$) in Plastic 1 Swamp using Pc 1-08 data for un-channelized fluxes.
Table 2a	Quarterly and annual inputs, outputs and retention of water discharge (m^3), major ions ($\text{meq m}^{-2} \text{ quarter}^{-1}$) and DOC ($\text{g m}^{-2} \text{ quarter}^{-1}$) in Plastic 1 Swamp using lysimeter data for un-channelized fluxes.
Table 2b	Quarterly and annual inputs, outputs and retention of major alkalinity components ($\text{meq m}^{-2} \text{ quarter}^{-1}$) in Plastic 1 Swamp using lysimeter data for un-channelized fluxes.
Table 3	Quarterly and annual inputs and outputs of discharge weighted concentrations of major ions ($\mu\text{eq/L}$) and DOC (mg/L) with the coefficient of variation (C.V.) as well as charge balance ($\mu\text{eq/L}$ and %) with the mean and standard deviation.

LIST OF FIGURES

- Figure 1 Location of Plastic 1 swamp, major inflows and outflow .
- Figure 2 Quarterly and annual input cation fluxes to Plastic 1 swamp.
- Figure 3 Quarterly and annual input anion fluxes to Plastic 1 swamp.
- Figure 4 Quarterly and annual discharge at Pc 1-08 inflow to Plastic 1 swamp.
- Figure 5 Quarterly and annual output cation fluxes to Plastic 1 swamp.
- Figure 6 Quarterly and annual output anion fluxes to Plastic 1 swamp.
- Figure 7 Relationship between the daily average summer precipitation depth and
number of days of the year with zero discharge from the Plastic 1 swamp.
- Figure 8 Quarterly and annual retention of cations in Plastic 1 swamp.
- Figure 9 Quarterly and annual retention of anions in Plastic 1 swamp.
- Figure 10 Quarterly and annual retention of alkalinity components in Plastic 1 swamp.

Figure 11 Quarterly and annual retention of computed alkalinity (sum of components in Figure 10) in Plastic 1 swamp.

INTRODUCTION

Minerotrophic conifer swamps are common peatlands on the Precambrian Shield in Canada, with important impacts on down-stream and lake chemistry (Gorham et al. 1984; Dillon & LaZerte 1992). Recently there has been a resurgence of interest in the impact of peatlands on water chemistry (Gorham et al. 1984; Bayley et al. 1986; Bayley et al. 1987; DeVito et al. 1989; Urban et al. 1989a; DeVito & Dillon 1992). Traditional minerotrophic swamp inputs are non-acidic, however when the surrounding uplands have been anthropogenically acidified, elevated inputs of protons, aluminum and sulphate are introduced into the swamp. How peatlands transform and/or retain these compounds and their impact on other peatland processes is an important subject of study (Gorham et al. 1984). Also of interest is how climatic conditions, in particular the amount of rainfall, affect this transformation and/or retention in peatlands (Gorham et al. 1984).

Detailed long-term chemistry and input-output budgets are provided here for a representative minerotrophic conifer swamp (Pc 1) on the PreCambrian Shield in the Muskoka-Haliburton area of Ontario. Previous studies of this swamp have examined the hydrology (Shibitani 1988; Wels & DeVito 1989), paleoecology (Warner et al. 1992), the transformation of nutrients from inorganic to organic forms (DeVito et al. 1989; DeVito & Dillon 1992) and the retention of as well as transformation to organic form of some metals (Al, Pb, Cu; LaZerte 1989; LaZerte et al. 1989; LaZerte 1991). Kirchner et al. (1992; Figure 1) have described the effect of drought on the swamp's outflow chemistry in the

context of testing a geochemical model of stream chemistry. The hydrology and geochemistry of the surrounding uplands have also been analyzed (Wels et al. 1991; Kirkwood & Nesbitt 1991; Law 1991; Dillon & LaZerte 1992). DeVito and Dillon (1992) examine the role of hydrology on phosphorus and nitrogen dynamics in the Pc 1 conifer swamp. In this study, the role of hydrology (low swamp water level in particular) and upland acidification on the net retention of the base cations, sulphate, aluminum species and alkalinity is determined.

METHODS

The Muskoka-Haliburton region is located on a southern extension of the Pre-Cambrian Shield in Central Ontario. The sensitivity of this area to acidic deposition, coupled with the relatively high historical deposition of acidity have made the Muskoka-Haliburton region one of the most acidified in Ontario (Neary & Dillon 1988). Precipitation depth in the Muskoka-Haliburton area averages about 1 m/yr (Dillon et al. 1988) with maximum precipitation depth usually in the fall (September to November) and minimum in the summer (June to August). Winter (December to February) precipitation accumulates as snow pack until spring (March-May) snowmelt.

The interior of the three largest basins draining into Plastic Lake is dominated by conifer swamps, of which the Plastic 1 (Pc 1) swamp and sub-catchment are the largest (Figure 1). The upland soils of the Pc 1 catchment surrounding the swamp are sandy, shallow (~0.5 m)

podzols underlain by granitic gneiss and amphibolite (Lozano 1987; Kirkwood & Nesbitt 1991; Law 1991). The upland forest cover is dominated by white pine, hemlock and soft maple. The conifer swamp soils are peaty, organic mucks and gleysols (mean depth 2-3 m; 7 m max. depth) dominated by white cedar and black spruce with sphagnum sp. as the dominant ground cover (Lozano & Parton 1986; DeVito 1989).

For the purposes of this study, the Pc 1 basin above the swamp outflow (Pc 1-03) has been subdivided into three different sub-basins draining into the swamp itself (Figure 1). The surveyed areas (ha) are:

Pc 1-08	3.22
Pc 1-04	5.29
Other uplands	9.46
<u>Swamp</u>	<u>2.12</u>
Pc 1-03 (Total)	20.09

Of the three areas that drain into the swamp, Pc 1-08 is unique in that it has a fairly long, steadily descending reach with few ponding or wetland areas. The underlying bedrock forms a narrow notch just before entering the conifer swamp which concentrates the water flux and is the weir location. Of all the surface inflows to the swamp, Pc 1-08 discharges the greatest amount over the longest period of time. Chemical data from Pc 1-08 is available from March 1983. Discharge data is available from December 1986; prior to that, discharge must

be estimated by areal pro-rating discharge at Pc 1, a weir 50 m downstream from the swamp outflow.

The Pc 1-04 sub-basin is larger than Pc 1-08. The headwaters of this sub-basin are a small alder swamp that drains across the top of the catchment and cascades down into the main swamp at several locations. All possible sampling sites flow erratically and represent only a small portion of the total flow from this sub-basin. For chemical sampling, a site was chosen that appears to have the greatest flow but a weir was not installed. Hydrological discharge from this area is computed by areal pro-rating of the discharge from Pc 1-08 or Pc 1 (prior to December 1986 only). Chemical data from Pc 1-04 is available from March 1983.

The largest contributor to the Pc 1 conifer swamp is not channelized or surface flow at all, but macropore flow along the bedrock soil interface. The porous, sandy upland soils allow rapid vertical permeation of snowmelt and rainfall down to the bedrock interface. This is followed by gravitational, macropore flow along the bedrock soil/interface down to the swamp (Shibitani 1988; Wels et al. 1991). Areal pro-rating of the discharge from Pc 1-08 or Pc 1 (prior to December 1986 only) is used to estimate the hydrological discharge from these areas.

There are several possible estimates of the chemistry of the unchannelized macropore flow into the swamp:

- i) Samples collected at Pc 1-04 can be used. However, because of the headwaters swamp in Pc 1-04, these samples contain somewhat elevated amounts of dissolved organic carbon (DOC) relative to Pc 1-08 and ground water samples.
- ii) Pc 1-08 is more representative of groundwater flux than Pc 1-04, as its DOC levels are more typical of ground water samples. However, inorganic aluminum can drop significantly in the Pc 1-08 stream bed (LaZerte 1989).
- iii) Perhaps the best choice is to use ground water samples collected from 12 lysimeter pits installed in the northern section of the Pc 1 catchment (Figure 1). Although the pits are not randomly distributed about the ungauged portion of the catchment, their locations are representative of typical soil conditions in the catchment: there is as much variation in soil properties within the Pc 1-08 sub-basin as there is across all the upland soils of the Pc 1 catchment (Lozano 1987). As soil water chemistry changes dramatically with depth (LaZerte 1989), the samples used here are only those relatively uniform (over both time and space) samples collected deep in the mineral soil horizon (usually a Bf2 horizon) close to the bedrock interface. Only the four most recent years of data are available from these pits, so for longer term analysis (including all graphs) , Pc 1-08 data is used.

The final input to the Pc 1 conifer swamp is direct precipitation to its surface. Although this forms a relatively small proportion of all inputs (only 10.6% by area), it can be an important

contributor during the summer when other inputs are dry. A bulk deposition collector and meteorological site located in the Plastic Lake catchment provide weekly chemistry and daily precipitation depth (OME 1990). Data from this site is available from December 1983. Earlier data (March to December 1983) is obtained from a similar site located within 15 km. The bulk collectors used at these sites appear to give good estimates of both wet and dry deposition of sulphate (Dillon et al. 1988). Two empirical models were calibrated against three years of snowpack monitoring in Pc 1 catchment (Findeis et al. 1989) to account for the substantial amounts of deposition collected in the swamp snowpack over winter and not released until spring snowmelt. The first model uses average daily temperature and daily precipitation depth to estimate the accumulation and melt of snowpack water equivalents over winter to spring. The second model uses empirically estimated exponential decay constants for each major chemical parameter in the snowpack to estimate changes in concentration over time (LaZerte, unpubl. results). Taken together, the two models provide daily chemical fluxes from the snowpack directly onto the swamp.






The outflow from the Pc 1 conifer swamp is monitored chemically at a well defined bedrock lip (Pc 1-03) which controls the water level of the swamp. The hydrologic discharge at this site is areally pro-rated against the discharge at the Pc 1 weir located about 50 m downstream. Chemical data from Pc 1-03 and hydrological data from Pc 1 are available from March 1983.

The V-notch weirs at Pc 1 and Pc 1-08 are established on a bedrock base with little evidence of leakage. The notches are heated to eliminate ice buildup and provide year-round operation. The stage/discharge relations of the weirs are individually calibrated using Pygmy meters in standard reaches (high flow) or buckets (low flow). The stage is continuously monitored in a heated stilling well using a float connected to a Leopold-Stevens A71 recorder. Daily discharge is computed from the digitized strip charts.

Samples were collected for chemical analyses more frequently during high flow periods (1-4 times per week in the spring and fall) than during low flow periods (once per week if water was available). All data was carefully examined for outliers using time trend plots and charge balance calculations. Linear interpolation was used to estimate the chemistry on days between sample collection. Sampling was frequent enough and discharge-concentration relationships weak enough that filling in missing data by interpolation was more accurate than using discharge-concentration relationships.

Chemical analyses were performed as described in OME (1983). All analyses were insensitive to DOC interference, including the ion chromatographic sulphate method. Aluminum species were estimated with a catechol violet / anion exchange method (LaZerte et al. 1988) and with the calculations described in LaZerte (1984). Oliver's model (Oliver et al. 1983) was used to estimate the organic anion contribution to charge balance. The model works well in these waters and elsewhere when corrected for iron and aluminum complexation (Oliver et al. 1983; LaZerte and Dillon 1984; Cantrell et al. 1990).

Alkalinity was both titrated to a fixed endpoint of pH 3.8 and calculated at surface water sites. At groundwater sites, alkalinity was only calculated. For these calculations, alkalinity was defined to include bicarbonate alkalinity as well as the alkalinity generating portion of inorganic Al and weak organic acids. The latter was estimated as that fraction of Oliver's model organic acid (Oliver et al. 1983) that contributes to alkalinity titrations down to a pH of 3.8 (Orgw⁻). The formula for calculating this alkalinity follows from the proton balances of the individual acids that are as weak or weaker than the arbitrarily defined weak organic acids at pH 3.8 (Stumm and Morgan 1981):



etc.

which lead to :

$$\begin{aligned} (1) \text{ Alkalinity} = & [\text{Orgw}^-] + 2[\text{CO}_3^{2-}] + [\text{HCO}_3^-] + [\text{AlOH}^{2+}] + 2[\text{Al}(\text{OH})_2^+] + 3[\text{Al}(\text{OH})_3] \\ & + 4[\text{Al}(\text{OH})_4] + [\text{OH}^-] - [\text{H}^+] \end{aligned}$$

$\text{Al}(\text{OH})_3$ includes monomeric, polymeric and colloidal forms that respond to the titration (see also the Appendix).

The start of each hydrological year was taken to be 1 September, before the fall rains began and when discharge from the swamp was usually zero. Input and output chemical fluxes per square metre of swamp surface were prepared on a daily basis and then quarterly and annual sums computed. Input and output volume weighted concentrations on a quarterly basis were calculated by dividing quarterly chemical flux by the equivalent quarterly water flux and adjusting for the swamp area.

ERRORS and METHOD CHECKS

A regression (with no intercept) of the quarterly discharge of the Pc 1-08 inflow after December 1986 against the Pc 1 weir gives an R^2 of 0.99 and a slope of 0.16 ± 0.1 (2 S.E.). This slope compares favourably with the ratio of areas (0.15), confirming the use of areal pro-rating on a quarterly or annual basis for missing upland water fluxes. DeVito (1989) has also concluded that areal prorating works well in these catchments.

A comparison of the calculated alkalinity with an actual fixed end point alkalinity titration to pH 3.8 demonstrates an approximate 1:1 relationship ($R^2=0.81$). A multiple linear regression of titrated alkalinity versus the three major components (with no intercept) also reveals approximate 1:1 relationships for each component:

Variable	Coefficient	S.E.	P
$2[\text{CO}_3^{2-}] + [\text{HCO}_3^-] +$			
$[\text{OH}^-] - [\text{H}^+]$	0.94	0.02	0.0
$[\text{Orgw}^-]$	1.15	0.05	0.0
$[\text{AlOH}^{2+}] + 2[\text{Al}(\text{OH})_2^+] +$			
$3[\text{Al}(\text{OH})_3] + 4[\text{Al}(\text{OH})_4^-]$	1.06	0.05	0.0

The first term is the traditional definition of (bicarbonate) alkalinity; in all of these samples, its major component is $[\text{H}^+]$ (see also the Appendix).

The major systematic error in these Pc 1 swamp budgets is probably the estimate of groundwater chemical concentrations. A rough estimate of this error can be obtained by comparing the total input fluxes to the swamp using Pc 1-08 stream chemistry for unmonitored areas (Table 1a) with fluxes using lysimeter chemistry (Table 2a) over the same period. The following slopes were obtained regressing (with no intercept) the Pc 1-08 estimate against the lysimeter estimate using the most recent four years of quarterly data where both estimates are available:

Variable	Coefficient	S. E.	P
Base Cations	0.91	0.01	0.0
Inorganic Al	0.85	0.02	0.0
H ⁺	1.09	0.01	0.0
Sulphate	0.93	0.01	0.0
Orgw ⁻	0.90	0.04	0.0
Other Anions	0.82	0.01	0.0

With the exception of H⁺, the Pc 1-08 estimates of unchannelized groundwater chemistry are lower than estimates based on lysimeter data. The errors appear to be on the order of 8% lower for base cations and sulphate and 10-18% lower for the other ions.

Charge balances on the quarterly input and output volume weighted concentrations average $-7\% \pm 6\%$ (mean \pm standard deviation, n=30) and $-1\% \pm 6\%$ respectively, with maximum deviations of -24% and $+16\%$ respectively, for the long term budgets using Pc 1-08 chemistry for unchannelized input fluxes (Table 3).

RESULTS

Base cations are the major cations entering the swamp with positively charged aluminum species and H⁺ also important (Figure 2; Tables 1a, 2a). Sulphate dominates the anion inputs with Orgw⁻, bicarbonate, chloride and nitrate of lesser significance (Figure 3; Tables

1a, 2a). The major inputs occur in the spring during snowmelt, with fall and occasionally early winter (December) influxes being somewhat important.

As chemical flux is the product of chemical concentration and water flux, changes in chemical flux are caused by changes in either one or both. Changes in surface water flux is a major determinant of the temporal pattern of chemical input flux to the swamp (Figure 4; Table 3; quarterly C.V.=86%). The volume weighted input concentrations are comparatively stable (Table 3; quarterly C.V. are 23% and 18% for base cations and sulphate respectively). On an annual basis, both total cation inputs and total anion inputs have maximum and minimum annual fluxes (1984/85 and 1986/87 respectively) that correspond to maximum and minimum annual water fluxes (Figures 2-3).

Base cations and sulphate also dominate the swamp outflow, but H^+ is more important than inorganic aluminum cations and organic anions are more abundant (Figures 5-6; Tables 1a, 2a). Spring is very important to the output fluxes, but, in contrast to inputs and especially in the four years following the dry summers of 1983, 87, 88 and 89, Fall + Winter is even more important.

Summer drought conditions have a major impact on wetland chemical concentrations and consequently on chemical output fluxes. The average daily summer precipitation depth is correlated with the number of days with zero discharge from the Pc 1 swamp (Figure 7). A lowered water table and exposed muck soils are found during these periods of zero

discharge in the summer. In the years following an extended summer period of zero discharge, concentrations of most chemical parameters become elevated and, in particular, with the onset of fall rains the first flush of water from the swamp is very concentrated (Table 3; Figures 5-6; see also Kirchner et al. 1992, Figure 1). The variability of output water flux is similar to the inflow (Table 3; quarterly C.V.=88%), but the equally variable base cation and sulphate concentrations (Table 3; quarterly C.V.=68-82%) also have a substantial impact on the temporal variability of chemical fluxes from the swamp.

It is in the Fall+Winter quarters following the dry summers that the most substantial negative retentions (net export, Figures 8-9; Tables 1a, 2a) of base cations and sulphate occur. The annual pattern of sulphate and base cations is also revealing: only in the years following dry summers is there a net export of sulphate (negative retention). In all other years, including those years with the highest and lowest water fluxes through the swamp (1984/85 and 1986/87 respectively), sulphate is retained. Base cations are never positively retained on an annual basis (when using the Pc 1-08 estimate for ungauged area; Table 1a); they are approximately in balance after a normal summer and exported after drought, even through the highest and lowest water flux years. Clearly summer drought conditions have a substantial impact on the retention of these major ions, primarily by releasing large quantities of them in the subsequent fall and winter discharge periods.

The negative retention (net export) of H^+ shows an annual pattern that may have as much to do with annual water flux as summer drought. Large spring net exports dominate its

annual budget. Although the pH drops dramatically, most of the fall export of sulphate following summer drought is balanced by base cations, not protons. Nonetheless, the three years of minimum proton export (Figure 8) do correspond to the three years of positive sulphate retention (Figure 9). The seasonal and annual retentions of aluminum cations, Orgw^- and other anions show less pattern, although the three latest drought years (Figure 8) do have elevated aluminum retentions relative to the three previous wet years.

H^+ and aluminum hydroxide species are the two dominant components of alkalinity retention in the Pc 1 conifer swamp and consequently the temporal pattern of alkalinity retention is similar to theirs (Figure 10-11; Tables 1b, 2b). Annually, alkalinity is retained; only in the summer are small amounts of alkalinity exported (Figure 11; Tables 1b, 2b). This is the result of H^+ retained from summer rainfalls and Orgw^- export. If the contribution of aluminum hydroxides and weak organics to alkalinity were ignored, the estimates of alkalinity retained by the swamp would be less but the pattern would be similar.

Net DOC fluxes from this peatland range from 21 to 43 g $\text{m}^{-2} \text{yr}^{-1}$ (Tables 1a, 2a).

DISCUSSION

One of the most significant results of this research is the net export of base cations from the Pc 1 swamp over the current seven year study period. Other budget studies, although rare, indicate that both ombrotrophic and minerotrophic peatlands retain base cations (Hemond

1980; Gorham et al. 1984). The lack of base cation retention at this acidified site may be the result of competition for new exchange sites by incoming protons and aluminum ions. In the past, there may have been sufficient incoming base cations (and fewer competing protons and Al ions) to populate many of the sites. Under current conditions, however, with the loss of base cations and the increase of aluminum ions, the cation exchange sites in the swamp are likely becoming more acidified as predicted by Gorham et al. (1984) and Bayley et al. (1986).

Another important result is the greater retention of sulphate relative to base cations. Currently, the swamp retains sulphate during normal and wet summers, and then exports it after dry summers. A similar pattern has been observed in this swamp for phosphorus and nitrogen (Devito & Dillon 1992). It is likely that the exposure and oxidation of reduced sulphur compounds is responsible for these chemical fluctuations (Gorham et al. 1984; Bayley et al. 1986; Bayley et al. 1987; Urban et al. 1989b; DeVito & Dillon 1992); DeVito (unpubl. results) has documented increased rates of sulphur mineralization as the result of summer drought in Pc 1 swamp. This pulsed export of sulphate from the Pc 1 swamp is mostly balanced by an export of cations, presumably via an exchange reaction between the proton from the sulphur oxidation reaction and the base cations on the peat exchange sites. The retention of sulphate after wet summers, however, is not balanced by a retention of cations but by their reduced net export and, in part, by decreased proton export and increased aluminum retention. Geochemically, this is consistent with sulphate reduction increasing peat pH which reduces proton export but increases aluminum precipitation or

absorption. Eventually, if current conditions persist, the exchangeable pool of base cations in the swamp will be depleted by the periodic export of sulphate and more protons and some aluminum ions will be exported to compensate. This will increase acidity export (alkalinity retention) primarily by directly increasing the proton export and indirectly reducing Orgw^- export.

Net sulphate retention has been observed in other peatlands as has post-drought release (Hemond 1980; Gorham et al. 1984 and references therein; Bayley et al. 1986; Urban et al. 1989b), but no investigator, to my knowledge, has reported an annual net export of sulphate (or of base cations) from a minerotrophic peatland. Bayley et al. (1986) hypothesize that, other conditions being equal, peatlands with historically less anthropogenic sulphur deposition should currently have greater base cation neutralization capacity and hence less proton export coupled with sulphate export. Their results and this study both support this hypothesis.

The retention of alkalinity observed in this study is not unexpected. Urban et al. (1989a) have documented the role of peatland organic acid exports in the acidification of stream water. In the Pc 1 swamp, organic acids increase the acidity of already acidified water passing through the swamp (LaZerte and Dillon 1984). This is especially the case in wet years (e.g., 87/88; Tables 1a, 2a) when the relative proportion of organic anions to sulphate anions in the swamp outflow approaches one. Conversely, after dry summers the large release of sulphate from the swamp reduces the relative importance of organic acids. More

generally, the increased acidity of water passing through the swamp may be the result of high water yield (4-7 m of water m^{-2} of swamp yr^{-1}) which increases organic acid leaching from many peatlands, although the total and net flux (21 to 43 $\text{g m}^{-2} \text{yr}^{-1}$) of organic carbon from the wetland is not unusual (Urban et al. 1989a). Another likely cause is the gradual loss of exchangeable base cations from the Pc 1 conifer swamp; these base cations would neutralize some of the internally generated organic acidity as well as mineral acidity from sulphur oxidation.

More detailed study is required to determine the exact layers of peat involved in the cycle of sulphate accumulation and export, and whether these same layers are also responsible for the cation export. Presumably reduced sulphur is immobile and can only be exported when swamp water levels drop to the point where oxygen can diffuse to its location. However, the protons released by sulphur oxidation are mobile, and it may be that the depth of peat available for proton/cation exchange is greater than that available for the sulphur oxidation. In addition, possible internal sources of base cations diffusing up from deep sediments (as has been postulated for iron; LaZerte 1991) should be considered. Similarly, the role of different regions of the swamp (e.g., the Pc 1-08 lagg on the eastern side of the swamp; see DeVito & Dillon 1992) should be examined. The crucial question is the amount of residual sulphur available and the size of cation storage in the swamp available to neutralize these pulsed sulphur oxidation events.

Climatic changes in temperature and rainfall have been shown to influence lake chemistry in unexpected ways (Webster et al. 1990; Psenner and Schmidt 1992). With respect to peatland behaviour, it may nonetheless be possible to extrapolate the effects of the drought conditions observed in this and other studies (e.g., Bayley et al. 1986) to wetland behaviour under the longer-term reduced rainfall conditions predicted for some regions. It should be emphasized that only long-term studies can provide the perspective and data needed for long term predictions.

REFERENCES

- Bayley, S.E., D.H. Vitt, R.W. Newbury, K.G. Beaty, R. Behr and C. Miller. 1987. Experimental acidification of a sphagnum dominated peatland: first year results. *Can. J. Fish. Aquat. Sci.* 44: 194-205.
- Bayley, S.E., R.S. Behr, and C.A. Kelly. 1986. Retention and release of S from a freshwater wetland. *Wat. Air Soil Pollut.* 31: 101-114.
- Cantrell, K.J., S.M. Serkiz and E.M. Perdue. 1990. Evaluation of acid neutralizing capacity data for solutions containing natural organic acids. *Geochim. Cosmochim. Acta.* 54: 1247-1254.
- DeVito, K.J. 1989. Waterborne phosphorus and nitrogen retention in some Precambrian Shield wetlands: the role of hydrology and oxidation-reduction potential. M.Sc. Thesis, Trent University, Peterborough, Ontario, Canada. 256 pp.
- DeVito, K.J., P.J. Dillon and B.D. LaZerte. 1989. Phosphorus and nitrogen retention in five Precambrian Shield wetlands. *Biogeochem.* 8: 185-204.
- DeVito, K.J. and P.J. Dillon. 1992. The influence of hydrologic fluctuations and peat oxia on the phosphorus and nitrogen dynamics of a conifer swamp. Submitted.

- Dillon, P.J, M. Lusi, R. Reid and D. Yap. 1988. Ten-year trends in sulphate, nitrate and hydrogen deposition in Central Ontario. *Atmos. Envir.* 22: 901-905.
- Dillon, P.J. and B.D. LaZerte. 1992. Response of Plastic Lake catchment, Ontario, to reduced sulphur deposition. *Envir. Pollut.* 77: 211-217.
- Findeis, J.G., B.D. LaZerte and L.D. Scott. 1989. Biogeochemistry project: snow cores 1988-1989, Plastic Lake, Harp Lake, Hawkeye Lake. Report BGC-031, Dorset Research Centre, Ont. Min. Envir., Dorset, Ontario, Canada.
- Gorham, E., S.E. Bayley and D.W. Schindler. 1984. Ecological effects of acid deposition upon peatlands: a neglected field in "Acid-Rain" research. *Can. J. Fish. Aquat. Sci.* 41: 1256-1268.
- Hemond, H.F. 1980. Biogeochemistry of Thoreau's Bog, Concord, Massachusetts. *Ecol. Monogr.* 50: 507-526.
- Hemond, H.F. 1990. Acid neutralizing capacity, alkalinity, and acid-base status of natural waters containing organic acids. *Envir. Sci. Technol.* 24: 1486-1489.
- Kirchner, J.W., P.J. Dillon and B.D. LaZerte. 1992. Predicted response of stream chemistry to acid loading tested in Canadian catchments. *Nature.* 358: 478-482.

- Kirkwood, D.E. and H.W. Nesbitt. 1991. Formation and evolution of soils from an acidified watershed: Plastic Lake, Ontario, Canada. *Geochim. Cosmochim. Acta.* 55: 1295-1308.
- Law, K.R. 1991. The weathering of granitic tills and the development of soil profiles at Plastic and Harp Lake, Ontario. M.Sc. Thesis, University of Western Ontario, London, Ontario, Canada. 201 pp.
- LaZerte, B.D. 1984. Forms of aqueous aluminum in acidified catchments of Central Ontario: a methodological analysis. *Can. J. Fish. Aquat. Sci.* 41: 766-776.
- LaZerte, B.D. and P.J. Dillon. 1984. Relative importance of anthropogenic versus natural sources of acidity in lakes and streams of Central Ontario. *Can. J. Fish. Aquat. Sci.* 41: 1664-1677.
- LaZerte, B.D. 1989. Aluminum speciation and organic carbon in waters of Central Ontario. In: Lewis, T.E. (ed.). *Environmental chemistry and toxicology of aluminum* (pp. 195-207). Lewis Publ. Chelsea, Michigan, USA.
- LaZerte, B.D., C. Chun and D. Evans. 1988. Measurement of aqueous aluminum species: comparison of dialysis and ion-exchange techniques. *Envir. Sci. Technol.* 22: 1106-1108.

- LaZerte, B.D., D. Evans and P. Grauds. 1989. Deposition and transport of trace metals in an acidified catchment of Central Ontario. *Sci. Tot. Env.* 87/88: 209-221.
- LaZerte, B.D. 1991. Metal transport and retention: the role of dissolved organic carbon. *Proc. 3rd Internat. Nordic Symp. on Humic Substances*, 21-23 August 1991, Turku, Finland. *Finnish Humus News*. 3: 71-82.
- Lozano, F, and W.J. Parton. 1986. Forest cover characteristics of the Harp #4 and Plastic #1 subcatchments at the southern biogeochemical study site. Report BGC-008, Dorset Research Centre, Ont. Min. Envir., Dorset, Ontario, Canada. 133 pp.
- Lozano, F. 1987. Physical and chemical properties of the soils at the southern biogeochemical study site. Report BGC-018, Dorset Research Centre, Ont. Min. Envir., Dorset, Ontario, Canada. 76 pp.
- Morel, F.M.M. 1983. Principles of aquatic chemistry. J. Wiley & Sons, New York. 446 pp.
- Neary, B.P. and P.J. Dillon. 1988. Effects of sulphur deposition on lake-water chemistry in Ontario, Canada. *Nature*. 333: 340-343.

Oliver, B.G., E.M. Thurman and R.L. Malcolm. 1983. The contribution of humic substances to the acidity of colored natural waters. *Geochim. Cosmochim. Acta.* 47: 2031-2035.

Ontario Ministry of the Environment. 1983. Handbook of analytical methods for environmental samples. Ont. Min. Envir., Rexdale, Ontario, Canada. 246 pp.

Ontario Ministry of the Environment. 1990. Meteorologic data procedures manual for the Dorset Research Centre. Ont. Min. Envir., Dorset, Ontario, Canada.

Psenner, R. and R. Schmidt. 1992. Climate driven pH control of remote Alpine lakes and effects of acid deposition. *Nature.* 356: 781-783.

Reuss, J.O. and D.W. Johnson. 1986. Acid deposition and the acidification of soils and waters. Springer-Verlag, New York. 119 pp.

Shibitani, R. 1988. Meltwater processes and runoff mechanisms in a small Precambrian Shield watershed during snowmelt. M.Sc. Thesis, Trent University, Peterborough, Ontario, Canada. 194 pp.

Stumm, W. and J.J. Morgan. 1981. Aquatic chemistry (2nd ed.). J. Wiley & Sons, New York. 780 pp.

- Urban, N.R., S.E. Bayley and S.J. Eisenreich. 1989a. Export of dissolved organic carbon and acidity from peatlands. *Wat. Resour. Res.* 25: 1619-1628.
- Urban, N.R., S.J. Eisenreich and D.F. Grigal. 1989b. Sulfur cycling in a forested sphagnum bog in northern Minnesota. *Biogeochem.* 7: 88-109.
- Warner, B.G., J.A. Janssens and S.G. Marsters. 1992. Paleoenvironmental history of a wetland near Plastic Lake, Dorset, Ontario. Report BGC-047, Dorset Research Centre, Ont. Min. Envir., Dorset, Ontario, Canada.
- Webster, K.E., A.D. Newell, L.A. Baker and P.L. Brezonik. 1990. Climatically induced rapid acidification of a softwater seepage lake. *Nature.* 347: 374-375.
- Wels, C. and K.J. DeVito. 1989. Tracing flow paths in a conifer swamp. In: Bardecki, M.J. and N. Patterson (eds.). Ontario wetlands: inertia or momentum (pp. 293-299) Federation of Ontario Naturalists, Don Mills, Ontario, Canada.
- Wels, C., C.H. Taylor, R.J. Cornett and B.D. LaZerte. 1991. Streamflow generation in a headwater basin on the Precambrian Shield. *Hydrol. Proc.* 5: 185-199.

APPENDIX - ALKALINITY METHOD

In alkalinity as defined here, the fixed pH 3.8 endpoint is arbitrary, but it does happen to be the pH in the Oliver model (Oliver et al. 1983) where 50% of the organic acids are dissociated and also where all carbonates are protonated and inorganic aluminum is not hydrolysed. Both Cantrell et al. (1990) and Urban et al. (1989a) recommend such a fixed endpoint alkalinity titration when organic acids and aluminum hydrolysis species are important. Hemond (1990) has presented an alternative approach that uses a simple correction factor applied to Gran alkalinity titrations in the presence of DOC; but it is unclear how titratable aluminum species would be corrected for.

Setting the reference proton level for the alkalinity titration to this level (3.8) implies that the dissociation of carbonic acid and HOrg_w and the hydrolysis of $\text{Al}(\text{H}_2\text{O})_6^{3+}$ are alkalinity neutral: their addition to or subtraction from solution does not change the alkalinity (Stumm and Morgan 1981) although the pH may change. However, the dissolution of calcium bicarbonate and aluminum trihydroxide do increase alkalinity by an amount equivalent to the introduction.

Others have defined alkalinity differently (e.g., Reuss and Johnson 1986; Hemond 1990) :

$$(2) \text{ Alkalinity} = C_B - C_A \text{ (Sum strong base cations - Sum strong acid anions)}$$

Then using charge balance, they obtain:

$$(3) \text{ Alkalinity} = [\text{Org}^-] + 2[\text{CO}_3^{2-}] + [\text{HCO}_3^-] + [\text{Al}(\text{OH})_4^-] + [\text{OH}^-] - [\text{H}^+] - 3[\text{Al}^{3+}] - 2[\text{AlOH}^{2+}] - [\text{Al}(\text{OH})_2^+]$$

where $[\text{Org}^-]$ is the concentration of ALL dissociated organic acids, both weak and strong as defined above. Thus, the dissociation of the strong organic acids at pHs less than 3.8 do not reduce alkalinity in this formulation (3) whereas they would in (1). Also, it is the dissolution of $\text{Al}(\text{OH})_3$ that is alkalinity neutral, not the hydrolysis of Al^{3+} . Although this definition (2) is close to (1) for bicarbonate waters, it does not agree well with the usual "alkalinity" titration with strong acid in the laboratory or in the field when inorganic aluminum, organic acids or CaCO_3 are abundant. Even though the organic acid discrepancy can be corrected by including the strong acid portion in C_A , a more difficult problem with the charge balance approach (2) is that it does not account for the weak acidic hydrolysis of positively charged metals (e.g., Al^{3+}), nor does it handle any uncharged species that may contribute to alkalinity such as CaCO_3 or $\text{Al}(\text{OH})_3$. As noted by Morel (1983):

"Although convenient, the conceptualization of alkalinity as a charge balance is not strictly correct; the proper concept is that of an acid-base balance."

Table 1a

DISCHARGE (m3), ION FLUXES (meq/m2), DOC flux (g/m2) Period	Discharge			Base Cations			Protons			Aluminum			Sulphate			Organic Anions			Other Anions			DOC		
	IN			IN			IN			IN			IN			IN			IN			IN		
	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET
Spring 83	76618	68513	8105	462	399	63	101	160	-59	38	3	34	595	467	128	8	85	-77	52	19	33	9	19	-10
Summer 83	5876	2980	2897	26	14	12	8	7	2	1	0	1	36	14	22	1	8	-7	6	1	5	0	2	-1
Fall 83	31718	25078	6639	184	458	-274	48	105	-57	24	12	12	275	510	-235	10	72	-62	41	35	5	4	15	-11
Win 83/84	25775	23918	1857	179	226	-47	47	82	-35	16	4	11	233	246	-13	5	42	-37	30	22	8	4	10	-7
Spring 84	69874	63516	6359	444	432	12	85	153	-68	50	8	42	536	445	91	20	103	-43	44	22	22	10	22	-12
Summer 84	13812	8889	4923	67	49	18	23	20	4	4	2	2	80	18	62	2	49	-47	15	6	9	2	9	-8
Fall 84	19451	13625	5826	109	146	-37	35	44	-9	12	3	9	137	111	25	4	47	-44	24	19	4	2	10	-8
Win 84/85	27175	26470	705	196	223	-27	36	74	-38	24	13	11	247	235	12	7	59	-52	22	17	5	4	11	-7
Spring 85	92142	84530	7612	540	515	25	119	207	-87	55	11	44	694	565	129	26	112	-66	60	38	22	12	25	-14
Summer 85	11597	6546	5051	56	30	26	53	15	8	3	1	2	68	4	64	2	38	-36	17	8	9	1	7	-6
Fall 85	39167	32173	6994	219	224	-5	51	83	-32	33	6	27	260	158	102	15	106	-90	32	32	0	5	21	-16
Win 85/86	11182	12986	-1804	74	92	-18	14	36	-22	13	3	10	88	95	-7	5	26	-21	9	12	3	2	5	-4
Spring 86	56282	56452	-170	346	348	-1	83	146	-63	61	11	50	412	370	42	29	100	-71	51	22	30	9	21	-12
Summer 86	15132	12294	2838	57	68	-11	27	30	-2	7	2	4	76	11	65	5	64	-58	20	11	9	2	12	-11
Fall 86	27811	34058	-6246	136	166	-31	35	73	-38	19	2	17	158	96	61	15	121	-106	33	25	8	5	23	-18
Win 86/87	5466	8011	-2545	36	39	-3	5	16	-11	4	0	4	40	30	10	2	19	-17	5	6	-1	1	4	-3
Spring 87	42533	41898	635	237	228	10	59	98	-39	44	8	36	294	241	53	23	70	-48	36	18	18	7	15	-8
Summer 87	5180	1735	3445	15	7	8	15	3	11	1	0	1	25	2	23	0	8	-8	10	2	8	0	1	-1
Fall 87	7093	3626	3467	38	175	-138	17	24	-7	6	6	0	49	195	-146	2	12	-10	16	12	3	1	3	-2
Win 87/88	23523	23417	106	202	394	-193	35	96	-61	41	13	28	229	432	-203	12	54	-42	40	37	4	3	12	-8
Spring 88	73722	64461	9261	530	520	10	93	177	-83	89	20	68	623	565	58	38	102	-64	80	54	26	11	22	-11
Summer 88	3784	772	3012	11	5	6	11	2	9	1	0	1	21	3	18	0	3	-3	7	0	7	0	1	-0
Fall 88	32876	23748	9129	220	377	-157	51	105	-54	48	12	36	273	384	-112	16	77	-61	51	38	13	5	16	-12
Win 88/89	6495	9001	-2506	48	106	-57	9	32	-23	11	4	7	58	107	-49	3	20	-17	11	15	-4	1	4	-3
Spring 89	72386	68263	4123	481	515	-34	94	194	-100	106	13	93	599	568	30	32	113	-81	73	52	21	10	25	-14
Summer 89	6356	4022	2334	28	22	6	11	10	1	4	1	3	39	10	28	2	22	-21	9	2	7	1	4	-3
Fall 89	31358	34958	-3600	190	516	-326	48	141	-93	41	19	22	218	491	-273	17	125	-108	51	56	-5	5	25	-21
Win 89/90	13992	16085	-2093	102	168	-66	23	57	-34	24	6	18	119	173	-54	7	35	-27	23	19	3	2	7	-5
Spring 90	66408	61379	5030	412	442	-29	79	158	-79	91	15	76	491	446	45	27	106	-80	65	46	19	9	22	-14
Summer 90	4911	2938	1973	18	20	-1	12	7	5	3	0	2	28	18	10	1	9	-8	7	1	6	0	2	-1
83/84	141180	121401	19778	873	1165	-291	203	360	-156	93	26	68	1125	1219	-94	37	267	-230	130	86	44	19	57	-37
84/85	150364	131170	19194	901	914	-13	213	339	-126	94	28	65	1146	916	231	39	257	-218	122	82	40	17	53	-34
85/86	121763	113905	7858	697	732	-35	175	295	-120	114	22	92	836	634	202	55	296	-241	112	76	36	17	60	-43
86/87	80990	85702	-4711	425	441	-16	113	190	-78	68	10	58	518	370	148	40	218	-179	84	51	33	13	43	-30
87/88	108121	92275	15846	780	1094	-314	156	299	-143	137	39	98	922	1195	-273	52	171	-120	144	104	40	15	37	-21
88/89	118113	105034	13079	777	1020	-243	165	341	-176	169	30	139	969	1070	-102	53	232	-180	143	106	37	17	49	-33
89/90	116659	115360	1309	723	1145	-422	162	363	-201	159	40	119	856	1128	-272	52	275	-223	146	123	23	16	56	-41

Table 1b

ALKALINITY FLUXES (meq/m ²)															
Period	Protons			Weak Organics			Aluminum			Bicarbonate			Total Alkalinity		
	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET
Spring 83	101	160	-59	8	20	-12	46	0	46	15	6	10	-31	-134	103
Summer 83	8	7	2	0	2	-2	2	0	2	1	0	1	-4	-5	0
Fall 83	48	105	-57	4	8	-5	34	0	34	6	2	5	-4	-94	90
Win 83/84	47	82	-35	4	8	-4	13	0	13	5	3	2	-26	-71	46
Spring 84	85	153	-68	10	23	-13	37	1	36	12	7	5	-27	-123	97
Summer 84	23	20	4	1	12	-10	6	0	5	4	1	3	-13	-7	-5
Fall 84	35	44	-9	2	9	-7	8	0	8	4	1	3	-21	-34	13
Win 84/85	36	74	-38	3	10	-7	18	0	18	6	2	4	-8	-61	52
Spring 85	119	207	-87	11	25	-14	89	1	88	16	8	8	-3	-172	169
Summer 85	23	15	8	1	9	-8	5	0	5	4	1	4	-13	-5	-8
Fall 85	51	83	-32	5	23	-17	29	0	29	10	2	9	-7	-59	52
Win 85/86	14	36	-22	2	5	-4	3	0	3	4	1	3	-5	-30	24
Spring 86	83	146	-63	9	21	-12	9	1	8	13	4	9	-52	-120	68
Summer 86	27	30	-2	2	15	-13	1	0	1	5	1	4	-19	-14	-5
Fall 86	35	73	-38	5	29	-24	8	0	8	13	3	10	-9	-41	31
Win 86/87	5	16	-11	1	5	-4	3	0	3	3	1	2	2	-10	12
Spring 87	59	98	-39	8	16	-9	20	0	20	8	4	4	-23	-78	55
Summer 87	15	3	11	0	2	-2	1	0	1	1	0	1	-12	-1	-11
Fall 87	17	24	-7	0	0	0	2	0	2	1	0	1	-13	-23	10
Win 87/88	35	96	-61	4	6	-3	11	0	10	5	1	3	-16	-88	72
Spring 88	93	177	-83	12	20	-8	45	1	44	12	6	6	-25	-150	126
Summer 88	11	2	9	0	1	-1	1	0	1	1	0	1	-10	-2	-8
Fall 88	51	105	-54	5	8	-3	17	0	17	6	1	5	-23	-95	72
Win 88/89	9	32	-23	1	3	-2	4	0	4	2	1	1	-3	-28	26
Spring 89	94	194	-100	11	22	-11	30	1	30	13	7	6	-40	-165	125
Summer 89	11	10	1	1	5	-4	2	0	2	2	0	2	-6	-5	-2
Fall 89	48	141	-93	4	15	-11	30	1	30	5	2	3	-9	-124	115
Win 89/90	23	57	-34	2	5	-3	9	0	9	4	2	2	-8	-50	42
Spring 90	79	158	-79	9	22	-14	44	1	43	14	6	7	-13	-128	115
Summer 90	12	7	5	0	2	-2	1	0	1	1	0	1	-10	-5	-5
83/84	203	360	-156	19	50	-32	89	1	88	27	12	15	-69	-296	227
84/85	213	339	-126	18	53	-35	120	1	119	30	12	18	-46	-273	227
85/86	175	295	-120	18	63	-46	42	1	41	32	8	24	-83	-222	139
86/87	113	190	-78	13	53	-39	32	1	32	25	7	18	-42	-130	88
87/88	156	299	-143	16	27	-11	59	1	57	18	7	11	-64	-263	200
88/89	165	341	-176	17	38	-21	53	1	52	23	9	14	-71	-292	221
89/90	162	363	-201	15	45	-30	84	1	83	23	10	13	-39	-307	267

Table 2a

DISCHARGE (m3), ION FLUXES (meq/m2), DOC flux (g/m2)	Period	DOC flux (g/m2)																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																
		Discharge						Base Cations						Protons						Aluminum						Sulphate						Organic Anions						Other Anions						DOC																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																						
		IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT

Table 2b

Period		ALKALINITY FLUXES (meq/m2)														
		Protons			Weak Organics			Aluminum			Bicarbonate			Total Alkalinity		
		IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET	IN	OUT	RET
Fall 86	33	73	-40	9	29	-21	1	0	1	18	3	15	-6	-41	35	
	Win 86/87	4	16	-12	1	5	-4	2	0	2	4	1	3	2	-10	12
Spring 87	55	98	-44	8	16	-9	29	0	29	13	4	9	-4	-78	73	
Summer 87	14	3	11	0	2	-2	1	0	1	1	0	1	-12	-1	-10	
Fall 87	17	24	-7	2	0	2	2	0	2	1	0	1	-11	-23	12	
Win 87/88	32	96	-64	6	6	-0	14	0	14	8	1	7	-4	-88	84	
Spring 88	86	177	-90	12	20	-8	65	1	64	21	6	16	11	-150	162	
Summer 88	11	2	9	0	1	-0	1	0	1	1	0	1	-10	-2	-8	
Fall 88	49	105	-55	8	8	-1	21	0	21	9	1	7	-12	-95	83	
Win 88/89	9	32	-23	1	3	-2	5	0	5	2	1	1	-1	-28	28	
Spring 89	86	194	-108	12	22	-10	55	1	55	25	7	19	7	-165	172	
Summer 89	11	10	1	1	5	-4	3	0	3	1	0	1	-6	-5	-1	
Fall 89	43	141	-99	6	15	-9	36	1	36	9	2	7	9	-124	133	
Win 89/90	21	57	-36	3	5	-3	11	0	11	4	2	3	-3	-50	47	
Spring 90	71	158	-87	9	22	-13	61	1	60	25	6	19	24	-128	152	
Summer 90	12	7	5	0	2	-2	2	0	2	1	0	1	-9	-5	-4	
86/87	106	190	-84	18	53	-34	33	1	32	36	7	28	-19	-130	110	
87/88	146	299	-153	21	27	-7	81	1	80	32	7	25	-13	-263	250	
88/89	155	341	-186	22	38	-16	85	1	84	37	9	28	-11	-292	281	
89/90	146	363	-217	18	45	-27	110	1	109	39	10	29	21	-307	328	

Table 3

Period	Volume weighted concentrations (µeq/L, mg/L for DOC)										Aluminum		Sulphate		Organic Anions		Other Anions		DOC		Charge Balance		% Charge Balance	
	Base Cations		Protons		IN		OUT		IN															
Spring 83	128	111	28	44	10	1	165	129	2	23	14	5	3	5	-15	-2	-9	-1						
Summer 83	94	51	30	25	3	0	130	51	4	30	21	2	2	6	-28	-6	-20	-8						
Fall 83	123	307	32	70	16	8	184	342	7	48	27	24	3	10	-47	-29	-24	-7						
Win 83/84	147	186	39	67	13	3	192	203	4	35	25	18	3	9	-23	1	-11	1						
Spring 84	135	131	26	47	15	2	163	135	6	31	13	7	3	7	-6	7	-4	4						
Summer 84	103	75	36	31	6	3	122	27	3	75	23	9	2	14	-3	-3	-2	-3						
Fall 84	119	159	38	49	13	3	149	122	4	52	26	21	3	11	-9	16	-5	8						
Win 84/85	153	174	28	58	19	10	193	184	5	46	17	13	3	8	-15	-1	-7	-0						
Spring 85	125	119	27	48	13	3	160	130	6	26	14	9	3	6	-15	4	-9	2						
Summer 85	102	55	42	27	5	1	125	8	3	70	31	14	2	13	-10	-9	-7	-10						
Fall 85	119	122	28	45	18	3	141	86	8	57	17	17	3	11	-2	10	-1	6						
Win 85/86	141	175	26	68	24	5	167	180	9	49	17	22	3	10	-2	-3	-1	-1						
Spring 86	131	131	31	55	23	4	155	140	11	38	19	8	3	8	-1	5	-0	2						
Summer 86	80	96	39	42	9	3	107	15	7	89	27	15	2	17	-14	21	-10	16						
Fall 86	104	127	27	56	14	1	120	73	11	92	25	19	3	18	-12	-1	-8	-1						
Win 86/87	140	153	18	62	17	2	156	118	7	74	20	23	3	15	-9	2	-5	1						
Spring 87	119	114	29	49	22	4	147	120	11	35	18	9	4	7	-6	-2	-4	1						
Summer 87	63	30	60	14	4	0	104	8	1	33	40	8	2	6	-18	-6	-13	-12						
Fall 87	113	525	51	72	18	17	148	584	5	36	47	37	2	8	-17	-43	-9	-7						
Win 87/88	182	356	32	87	37	12	206	390	11	49	36	33	3	11	-3	-3	-1	-4						
Spring 88	153	150	27	51	26	6	180	163	11	29	23	16	3	6	-9	-2	-4	-1						
Summer 88	63	27	62	12	6	1	117	18	2	18	42	2	2	3	-30	2	-20	5						
Fall 88	142	243	33	68	31	8	176	248	10	50	33	24	3	11	-13	-3	-6	-1						
Win 88/89	158	346	31	106	35	13	191	350	10	66	35	49	3	14	-12	0	-5	0						
Spring 89	141	151	28	57	31	4	176	167	10	33	21	15	3	7	-7	-3	-3	-2						
Summer 89	95	74	36	33	14	2	129	35	6	75	31	7	2	14	-21	-8	-14	-7						
Fall 89	129	350	33	96	28	13	148	333	11	84	35	38	3	17	-4	3	-2	1						
Win 89/90	155	255	34	86	36	9	181	263	11	53	34	30	3	11	-1	5	-0	1						
Spring 90	132	141	25	50	29	5	157	143	9	34	21	15	3	7	0	5	0	3						
Summer 90	80	85	52	30	12	2	119	76	4	40	32	6	2	7	-12	-5	-8	-4						
C.V	23	68	30	42	54	87	18	82	47	42	34	66	19	39	Mean	S.D.	10	12	6	6				
83/84	131	175	31	54	14	4	169	183	6	40	20	13	3	9	-18	-3	-10	-1						
84/85	127	129	30	48	13	4	162	129	5	36	17	12	3	8	-14	4	-8	2						
85/86	122	128	30	51	20	4	146	111	10	52	20	13	3	10	-3	8	-2	4						
86/87	111	116	30	50	18	3	136	97	10	57	22	13	3	11	-9	1	-6	0						
87/88	153	215	31	59	27	8	181	235	10	34	28	20	3	7	-9	-7	-4	-3						
88/89	140	183	30	61	30	5	174	192	10	42	26	19	3	9	-10	-3	-5	-1						
89/90	132	208	30	66	29	7	156	205	9	50	27	22	3	10	-2	4	-1	1						
C.V	10	25	2	12	33	39	10	32	25	20	18	27	6	17	Mean	S.D.	9	5	3	2				

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7

Plastic 1 Swamp





Figure 8



Quarter

Figure 9




Figure 10






Figure 11

Alkalinity Retention



Plastic 1 Swamp



Quarter

